

## QUESTION 1

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		


For the solution of this question please use only the front face and if necessary the back face of this page.

[5 pt] a) Graph the function of  $f(x) = e^{x+1} - 2$  by using horizontal and vertical shifts.

[10 pt] b)  $\lim_{x \rightarrow 4} \frac{(\sqrt{x^2 - 7} - x + 1) \sin(x - 4)}{(x^2 - 3x - 4)^2} = ?$  (Do not use the L'Hopital's Rule)

[10 pt] c) Let  $f(x)$  be the function illustrated in the figure below. For what values of  $x$  is  $f(x)$  discontinuous on  $[-4, 4]$ ? Classify the types of discontinuities. Give reasons for your answer.

c)




 $f$  is discontinuous at  $x = -2, 0$  and  $1$ .at  $x = -2$ :  $\lim_{x \rightarrow -2^-} f(x) = 2, \lim_{x \rightarrow -2^+} f(x) = 2$  $f$  has a removable discontinuity at  $x = -2$ at  $x = 0$ ,  $\lim_{x \rightarrow 0^-} f(x) = 0$ , but  $\lim_{x \rightarrow 0^+} f(x) = \infty$ , therefore, $f$  has an infinite discontinuity at  $x = 0$ at  $x = 1$ , since  $\lim_{x \rightarrow 1^-} f(x) = 1 \neq \lim_{x \rightarrow 1^+} f(x) = -1$ , $f$  has a jump discontinuity at  $x = 1$ .

b)  $\lim_{x \rightarrow 4} \frac{(\sqrt{x^2 - 7} - (x-1)) \sin(x-4)}{(x-4)^2 (x+1)^2} \stackrel{0}{=} \lim_{x \rightarrow 4} \frac{\sin(x-4)}{(x-4)} \cdot \lim_{x \rightarrow 4} \frac{[\sqrt{x^2 - 7} - (x-1)][\sqrt{x^2 - 7} + (x-1)]}{(x-4)(x+1)^2 [\sqrt{x^2 - 7} + (x-1)]}$

$$= \underbrace{\lim_{x \rightarrow 4} \frac{\sin(x-4)}{(x-4)}}_1 \cdot \underbrace{\lim_{x \rightarrow 4} \frac{x^2 - 7 - (x^2 - 2x + 1)}{(x-4)(x+1)^2 [\sqrt{x^2 - 7} + (x-1)]}}_{\frac{2}{(x-4)(x+1)^2 [\sqrt{x^2 - 7} + (x-1)]}} = \lim_{x \rightarrow 4} \frac{2}{(x-4)(x+1)^2 [\sqrt{x^2 - 7} + (x-1)]}$$

$$= \frac{2}{25.6} = \frac{1}{75.6}$$

a)  $f_1 = e^x$   
 $f_2 = e^{x+1}$   
 $f = e^{x+1} - 2$



## QUESTION 2

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		

For the solution of this question please use only the front face and if necessary the back face of this page.

[10 pt] a) Show that the function  $f(x) = \frac{1}{x}$  satisfies the Mean Value Theorem on the interval  $[a, b]$  such that  $a, b > 0$  and find the value of  $c$  in the conclusion of the theorem.

[15 pt] b) Find the equation of the normal line of the curve  $\sin(2xy) + y^2 - 2xy - \tan(\pi x) = 1$  at the point  $P(0, 1)$ .

a)  $\left\{ \begin{array}{l} \text{The Mean Value Theorem:} \\ \text{i) is continuous on } [a, b] \\ \text{ii) is differentiable in } (a, b) \\ \text{then, there is at least one point } c \text{ in } (a, b) \\ \text{at which } f'(c) = \frac{f(b) - f(a)}{b-a} \end{array} \right\}$

$f(x) = \frac{1}{x}, \quad a, b > 0$

i) Is  $f$  continuous on  $[a, b]$ ?  
 $f$  is undefined at  $x=0 \notin [a, b] \Rightarrow f$  is con

ii) Is  $f$  differentiable in  $(a, b)$ ?  
Yes,  $f'(x) = -\frac{1}{x^2}, x \neq 0$

There is at least one point  $c$  in  $(a, b)$ ;

$f'(c) = \frac{f(b) - f(a)}{b-a} \Rightarrow -\frac{1}{c^2} = \frac{\frac{1}{b} - \frac{1}{a}}{b-a} \Rightarrow -\frac{1}{c^2} = \frac{\frac{1}{b-a}}{\cancel{ab}(b-a)} \Rightarrow c^2 = ab$

$\cancel{a>b} \quad \cancel{ab(b-a)}$

$c = \sqrt{ab}$

$c \in (a, b), \quad 0 < a < b.$

b)  $\sin(2xy) + y^2 - 2xy - \tan(\pi x) = 1, \quad y = y(x)$

$$\Rightarrow (2xy)' \cos(2xy) + 2yy' - 2(xy)' - \pi \sec^2(\pi x) = 0$$

$$\Rightarrow 2[y+xy'] \cos(2xy) + 2yy' - 2[y+xy'] - \pi \sec^2(\pi x) = 0$$

$$\Rightarrow y' [2x \cos(2xy) + 2y - 2x] = \pi \sec^2(\pi x) - 2y \cos(2xy) + 2y$$

$$y' = \frac{\pi \sec^2(\pi x) - 2y \cos(2xy) + 2y}{2x \cos(2xy) + 2y - 2x} \Rightarrow m_{\text{tangent}} = y'|_{P(0,1)} = \frac{\pi \sec^2 0 - 2 \cos 0 + 2}{0 + 2 - 0} = \frac{\pi}{2}$$

$$m_{\text{normal}} = -\frac{1}{m_{\text{tangent}}} = -\frac{1}{\frac{\pi}{2}} = -\frac{2}{\pi}$$

The normal line equation:  $y - y_0 = m_{\text{normal}} (x - x_0)$

$$y - 1 = -\frac{2}{\pi} (x - 0) \Rightarrow y = -\frac{2}{\pi} x + 1$$

## QUESTION 3

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
Signature:	Electronic Post(e-mail) address:	Student Number:		

For the solution of this question please use only the front face and if necessary the back face of this page.

[8 pt] a) Let the values  $f(2) = \frac{\pi}{2}$ ,  $g(2) = -1$ ,  $f'(2) = 4$  and  $g'(2) = 0$  be given for two differentiable functions  $f(x)$  and  $g(x)$ . Find  $h'(2)$  for the function  $h(x) = 1 + \cos[f(x)g(x)]$ .

[17 pt] b) Find the asymptotes of the function (if any)  $f(x) = \frac{5x^2 + 8x - 3}{x^2 - 1}$ . Give reasons for your answer.

$$\begin{aligned} a) \quad h(x) &= 1 + \cos(f(x).g(x)) \Rightarrow \frac{dh(x)}{dx} = h'(x) = 0 + \frac{d}{dx}[f(x).g(x)].[-\sin(f(x).g(x))] \\ &\Rightarrow h'(x) = - (f'(x).g(x) + f(x).g'(x)).\sin(f(x).g(x)) \\ &\Rightarrow h'(2) = - (f'(2).g(2) + f(2).g'(2)).\sin(f(2).g(2)) \Rightarrow h'(2) = - (4.(-1) + \frac{\pi}{2}.0).\underbrace{\sin(\frac{\pi}{2}.(-1))}_{(-1)} \\ &= -4 // \end{aligned}$$

$$b) \quad f(x) = \frac{5x^2 + 8x - 3}{x^2 - 1} \quad f \text{ is not defined at } x = \pm 1.$$

For the vertical asymptotes:

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} \frac{5x^2 + 8x - 3}{(x-1)(x+1)} = \frac{5+8-3}{-0.2} = \frac{10}{-0} = -\infty \quad \boxed{x=1} \quad V.A.$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} \frac{5x^2 + 8x - 3}{(x-1)(x+1)} = \frac{5+8-3}{0.2} = \frac{10}{0} = \infty$$

$$\lim_{x \rightarrow -1^-} f(x) = \lim_{x \rightarrow -1^-} \frac{5x^2 + 8x - 3}{(x-1)(x+1)} = \frac{5-8-3}{-2.-0} = \frac{-6}{0} = -\infty \quad \boxed{x=-1} \quad V.A.$$

$$\lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow -1^+} \frac{5x^2 + 8x - 3}{(x-1)(x+1)} = \frac{5-8-3}{-2.0} = \frac{-6}{-0} = \infty$$

For the horizontal asymptotes: ( $\deg(5x^2 + 8x - 3) = \deg(x^2 - 1)$ )

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{5x^2 + 8x - 3}{x^2 - 1} = \lim_{x \rightarrow \infty} \frac{x^2(5 + \frac{8}{x} - \frac{3}{x^2})}{x^2(1 - \frac{1}{x^2})} = \frac{5}{1} = 5 \quad \boxed{y=5} \quad H.A.$$

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} \frac{(5 + \frac{8}{x} - \frac{3}{x^2})}{1 - \frac{1}{x^2}} = \frac{5}{1} = 5$$

No oblique asymptotes.

## QUESTION 4

The blanks below will be filled by students. (Except the score)

Surname:	Name:	Group Number:	List Number:	Score
		Student Number:		


For the solution of this question please use only the front face and if necessary the back face of this page.

[25 p] For the function  $f(x) = x\sqrt{4 - x^2}$ ,

- i) find the domain ~~and range~~
- ii) find the intervals on which the function is increasing and decreasing,
- iii) find the extrema and where they occur,
- iv) identify the concavity and, if any, find the points of inflection,
- v) sketch the graph.

i) its domain:  $4 - x^2 \geq 0 \Rightarrow |x| \leq 2$     its range: Since  $-2 \leq x \leq 2$ ,  $f(x) \in [-2, 2]$   
 $D = \{x \in \mathbb{R} \mid -2 \leq x \leq 2\}$

ii)  $f' = ?$      $f'(x) = 1 \cdot (4 - x^2) + x \cdot (-2x) = \frac{4 - x^2 - x^2}{(4 - x^2)^{1/2}} = \frac{4 - 2x^2}{(4 - x^2)^{1/2}} \Rightarrow f'(x) = \frac{2(2 - x^2)}{\sqrt{4 - x^2}}$ ,  
 $f'(x) = 0 \Rightarrow x = \pm \sqrt{2} \in D$



$f$  is decreasing on  $[-2, -\sqrt{2}] \cup (\sqrt{2}, 2]$   
 $f$  " increasing in  $[-\sqrt{2}, \sqrt{2}]$


iii) The critical points are  $x = -\sqrt{2}, \sqrt{2}$  ( $f'$  is zero) and  $f'$  changes its sign at these points.  
The end points are  $x = -2, 2$ .

$$f(-2) = 0 \quad f(-\sqrt{2}) = -2 \rightarrow \text{local minima (abs. minima)}$$

$$f(2) = 0 \quad f(\sqrt{2}) = 2 \rightarrow \text{local maxima (abs. maxima)}$$

iv)  $f'' = ?$      $f'' = \frac{d}{dx} \left[ \frac{2(2-x^2)}{\sqrt{4-x^2}} \right] = 2 \left\{ (-2x)(4-x^2)^{-1/2} - (2-x^2) \frac{(-2x)}{2\sqrt{4-x^2}} \right\} \frac{1}{(4-x^2)} = 2 \frac{x(x^2-6)}{(4-x^2)^{3/2}}$

$$f'' = 0 \Rightarrow x = 0 \in D, x = \pm \sqrt{6} \notin D$$



$f$  is concave up over  $(-\infty, 0)$   
 $f$  " down over  $(0, \infty)$

$f(0) = 0$  is the inf. point.

